



Mevet

Injectable formulations

Oral formulations
Medicated premixes
Feed supplements
Disinfectants

ENROVALL 50 mg/ml **Solution for injection** Enrofloxacin



Injectable formulations

Mevet

Polígono Industrial El Segre
Parc. 409-410
25191 Lleida (Spain)
T. +34 973 210 269
www.mevet.es

Marketed by

MPA Veterinary medicines and additives S.L.
Av. Corts Catalanes, 5-7
08173 Sant Cugat del Vallès (Spain)
T. +34 93 747 96 59
www.mpaveterinary.com

Injectable formulations

ENROVALL 50 mg/ml Solution for injection Enrofloxacin



COMPOSITION PER ml

Active substance:

Enrofloxacin 50 mg

Excipients:

Benzyl alcohol (E 1519) 14 mg

TARGET SPECIES

Cattle (calves) and pigs.

INDICATIONS FOR USE

Calves: treatment of infections of respiratory tract caused by strains of *Pasteurella multocida*, *Mannheimia haemolytica* and *Mycoplasma spp.* sensitive to enrofloxacin. Treatment of digestive tract infections caused by *Escherichia coli* strains sensitive to enrofloxacin. Treatment of sepsis caused by sensitive strains of *Escherichia coli* to enrofloxacin. Treatment of acute arthritis associated with mycoplasma, caused by strains of *Mycoplasma bovis* sensitive to enrofloxacin.

Pigs: treatment of infections of respiratory tract caused by strains of *Pasteurella multocida*, *Mycoplasma spp.* and *Actinobacillus pleuropneumoniae* sensitive to enrofloxacin. Treatment of digestive tract infections caused by *Escherichia coli* strains sensitive to enrofloxacin. Treatment of sepsis caused by sensitive strains of *Escherichia coli* to enrofloxacin.

WITHDRAWAL PERIOD

Calves: meat and offal: 12 days.

Not permitted for use in lactating animals producing milk for human consumption.

Pigs: meat and offal: 13 days.

DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION

Subcutaneous route in cattle and intramuscular in pigs.

Calves: 5 mg of enrofloxacin/kg b.w., corresponding to 1 ml of the product/10 kg b.w., once daily subcutaneously for 3 to 5 days. Acute arthritis associated with mycoplasma, caused by strains of *Mycoplasma bovis* sensitive to enrofloxacin: 5 mg of enrofloxacin/kg b.w., corresponding to 1 ml of the product/10 kg b.w. once a day subcutaneously for 5 days. Corresponding to 1 ml of product/10 kg b.w., once a day subcutaneously for 5 days.

Pigs: 2,5 mg of enrofloxacin/kg b.w., corresponding to 0,5 ml of the product/10 kg b.w., once a day intramuscularly for 3 days. In digestive tract infections or sepsis caused by *Escherichia coli*: 5 mg of enrofloxacin/kg b.w., corresponding to 1 ml of the product/10 kg b.w., once a day intramuscularly for 3 days.

SPECIAL PRECAUTIONS FOR STORAGE

Protect from light.

PACKAGE SIZES

Glass:

Box with 1 vial of 100 ml

Box with 1 vial of 250 ml

Box with 10 vials of 100 ml

Box with 15 vials of 250 ml

Polypropylene:

Box with 1 vial of 250 ml

Box with 15 vials of 250 ml

REGISTRATION No.

1131 ESP

