



Injectable formulations  
Oral formulations  
**Medicated premixes**  
Feed supplements  
Disinfectants

**LINCOVALL 55 g/kg**  
**Medicated premix**  
Lincomycin hydrochloride



**Medicated premixes**

**Mevet**  
Polígono Industrial El Segre  
Parc. 409-410  
25191 Lleida (Spain)  
T. +34 973 210 269  
[www.mevet.es](http://www.mevet.es)

Marketed by  
**MPA Veterinary medicines and additives S.L.**  
Av. Corts Catalanes, 5-7  
08173 Sant Cugat del Vallès (Spain)  
T. +34 93 747 96 59  
[www.mpaveterinary.com](http://www.mpaveterinary.com)

## Medicated premixes

### LINCOVALL 55 g/kg Medicated premix Lincomycin hydrochloride



#### COMPOSITION PER g

##### Active substance:

Lincomycin (hydrochloride) 55 mg

##### Excipients:

Almond – hazelnut shell flour

#### TARGET SPECIES

Pigs.

#### INDICATIONS FOR USE

**Pigs:** treatment and prevention of swine dysentery caused by *Brachyspira hyodysenteriae* sensitive to lincomycin. Treatment and prevention of enzootic pneumonia caused by *Mycoplasma pneumoniae* sensitive to lincomycin. The presence of the disease in the herd should be confirmed before preventive treatment.

#### WITHDRAWAL PERIOD

Meat and offal: 6 days.

#### DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION

In feed use.

**Dysentery:** 4,4 mg of lincomycin (hydrochloride)/kg b.w./day, for 3 weeks or until remission of symptoms (equivalent to 80 mg of premix/kg b.w./day).

**Enzootic pneumonia:** 8,8 mg of lincomycin (hydrochloride)/kg b.w./day for 3 weeks or until remission of symptoms (equivalent to 160 mg of premix/kg b.w./day).

Feed consumption depends on the clinical condition of the animal and the time of the year. To ensure a correct dosage, the concentration of premix in feed will be adjusted taking into account the daily consumption.

The weight of the animals should be determined as accurately as possible to avoid underdosing.

It is recommended to use the following formula:

$$\frac{\text{kg premix}}{\text{tonne of feed}} = \frac{\text{dose (mg/kg body weight)} \times \text{body weight (kg)}}{\text{daily intake (kg)} \times \text{premix dose (mg/g)}}$$

Example:

$$\frac{4 \text{ kg Lincovall 55}}{\text{tonne of feed}} = \frac{8,8 \text{ (mg/kg body weight)} \times 60 \text{ (kg)}}{2,4 \text{ (kg)} \times 55 \text{ (mg/g)}}$$

During granulation, it is recommended to maintain a temperature below 70 °C.

#### SPECIAL PRECAUTIONS FOR STORAGE

This veterinary medicinal product does not require any special storage conditions.

#### PACKAGE SIZES

Sack of 25 kg

#### REGISTRATION No.

258 ESP

